



PSYCHOLOGICAL CHALLENGES AND COPING MECHANISMS IN INDIVIDUALS WITH RHEUMATOID ARTHRITIS

Aksa Mughal^{*1}, Sikandar Hayat²

^{*1}PhD Scholar, Faculty of Medicine and Allied Health Sciences, The Islamia University of Bahawalpur, Pakistan; ²Assistant Professor, Faculty of Medicine and Allied Health Sciences, The Islamia University of Bahawalpur, Pakistan

^{*1}aksa_mughal43@gmail.com; ²yhayat@iub.edu.pk

Keywords

Rheumatoid arthritis, psychological well-being, mental health challenges, coping strategies, depression, anxiety, stress, loneliness, interpersonal difficulties, adaptive coping mechanisms

Article History

Received on 29 April 2024

Revised on 29 May 2024

Accepted on 15 June 2024

Published on 30 June 2024

Copyright ©Author

Corresponding Author: *

Abstract

This thesis explores the intricate relationship between mental health challenges and coping strategies in individuals with rheumatoid arthritis (RA). Beyond its physical symptoms, RA profoundly affects psychological well-being. The research investigates how mental health issues correlate with adaptive coping strategies in RA patients, using a correlational design and a sample of 200 patients (100 males and 100 females). The study employs standardized measures of depression, anxiety, stress, and loneliness to quantitatively assess participants. Findings reveal high rates of anxiety and depression among RA patients, with gender-specific differences: women reported more stress and anxiety, along with greater interpersonal difficulties, while men exhibited higher levels of depression. Women also demonstrated a wider range of coping strategies. Mental health issues were found to significantly predict coping mechanisms, underscoring the importance of psychological well-being in adapting to RA. These results highlight the need for personalized psychosocial interventions, tailored to the specific mental health needs of male and female RA patients. The study contributes to a deeper understanding of the psychological challenges faced by individuals with RA and calls for gender-sensitive treatments to enhance adaptive coping and overall mental health.

INTRODUCTION

The primary objective of this research is to investigate the prevalence of mental health concerns and explore the various coping strategies used by individuals diagnosed with rheumatoid arthritis (RA). The primary objective of this study is to examine and enhance understanding of the mental health difficulties encountered by persons who have rheumatoid arthritis (RA), while also exploring the coping mechanisms they use to manage

these issues. The primary objective of this research might involve a range of pursuits, including The main objective of this study is to evaluate the occurrence and characteristics of mental health issues, including depression, anxiety, and stress, as well as their level of severity, in persons who have been diagnosed with rheumatoid arthritis. Furthermore, this study aims to examine the coping mechanisms used by individuals with rheumatoid arthritis



in order to effectively address their mental health difficulties. This may include an examination of both adaptive and dysfunctional coping strategies.

According to the World Health Organization, chronic disease is identified as the primary factor contributing to early mortality on a global scale. Based on the estimations provided by the World Health Organization (WHO), it is accountable for 63% of all mortalities. Chronic disease is characterized by a gradual and prolonged course, necessitating patients to adjust to altered circumstances and impacting many facets of life, often resulting in adverse effects and a notable decline in health-related quality of life. Rheumatoid arthritis (RA) is a chronic condition that significantly reduces a person's functional capacity.


Rheumatoid arthritis (RA) is the prevailing rheumatic condition in the category of connective tissue disorders. The condition under discussion is characterized by a chronic and progressive inflammatory process originating from the synovial membrane. This process leads to the deterioration and destruction of articular tissues, resulting in impaired joint functionality. The age of onset for this condition often falls between 40 and 60 years, with a greater incidence seen in women compared to men, ranging from 3 to 4 times higher. Rheumatoid arthritis patients often experience a range of bodily manifestations, including joint deformities and degeneration, chronic pain, fatigue, unintended weight loss, and episodes of fever. Additionally, those who suffer from this condition face psychological difficulties that are typically characterised by negative emotions like anxiety, despair, a sense of bereavement, and interpersonal issues resulting from changes in the fulfilment of social obligations (Ziarko, 2014).

Rheumatoid arthritis (RA) is a medical condition that continues to have a significant impact on people's lives despite notable progress in its treatment (SCOTT et al., 2005). According to Symmons et al. (2005), at the individual level, there exists considerable heterogeneity in the extent of felt

impairment across individuals with similar levels of illness severity. Rheumatoid arthritis has a prevalence of around 400,000 among people aged 16 and above in the United Kingdom. The condition has the potential to impact individuals across all age groups. Typically, the development of this condition occurs during the age range of 30 to 50 years, which coincides with an individual's greatest period of productivity.

Rheumatoid arthritis (RA) is a chronic systemic autoimmune disease that mostly affects synovial joints, causing inflammation (synovitis), joint erosion, and cartilage degradation. Rheumatoid arthritis (RA) is a chronic systemic autoimmune disease that mostly affects synovial joints, causing inflammation (synovitis), joint erosion, and cartilage degradation. As a consequence, several patients experience a decline in functional status and an increase in disability. Rheumatoid arthritis (RA) may potentially present as extra-articular manifestations, impacting several organs in the body and resulting in increased rates of mortality and morbidity (Zielinski et al., 2019). Those who have rheumatoid arthritis may slowly develop a number of cardiovascular problems, such as pericarditis, myocarditis, pericardial and pleural effusions, congestive heart failure, and problems with their skin, eyes, gut, nervous system, or kidneys (Turesson, 2003).

The occurrence of rheumatoid arthritis (RA) is around 1% in the majority of nations. The number of new cases, also known as the incidence, has been documented as 40 per 100,000 individuals who meet the 2010 American College of Rheumatology (ACR)/EULAR categorization criteria at the beginning of the study (Humphreys et al., 2013). Even though strict disease management plans have been put in place, such as the "treat-to-target" (T2T) plan described by Smolen et al. (2017), Song et al. (2018) show that some rheumatoid arthritis (RA) patients still have high levels of disease activity. This evaluation considers tender and swollen joint counts, erythrocyte sedimentation rate, and visual analog core. The use of patients' Visual Analogue Scale and tender joint count as



factors within the scoring system (Song et al., 2018) illustrates one of the main drawbacks or benefits associated with mental health issues, which is the subjective nature of some elements. The subjective assessments used in this study may be susceptible to the impact of poor mood and depression among individuals with rheumatoid arthritis (Chancay et al., 2019).

Rheumatoid arthritis arises from a complex interaction between hereditary and environmental elements, including variables such as smoking and dietary preferences. The presence of a family member with the condition increases a person's susceptibility to it, even though the precise genetic mechanism underlying it is still unknown. Weight is an additional significant component that has been shown to have an elevated risk for those who are classified as overweight based on their body mass index (BMI) (Lempp et al., 2006). It is generally suggested for most individuals to have an optimal body mass index (BMI) falling between the range of 18.5 and 24.9. Deviating from this range by being underweight or obese may potentially increase an individual's vulnerability to certain health conditions. Moreover, the act of smoking substantially increases the likelihood of having rheumatoid arthritis. Dietary habits are also a contributing factor, since there is a positive correlation between excessive red meat consumption, inadequate vitamin C intake, and an elevated susceptibility to the illness. The combined influence of several variables has a role in the probability of acquiring rheumatoid arthritis (Lempp et al., 2006).

Rationale of the Study

The justification for doing a research investigation on the subject of mental health issues and coping mechanisms among persons diagnosed with rheumatoid arthritis (RA) is based on the significant influence that this enduring autoimmune condition has on the overall physical and emotional welfare of affected individuals. RA increases the risk of mental health issues like depression, anxiety, and stress. The presence of concurrent mental

health issues may have a substantial impact on the overall well-being of those diagnosed with rheumatoid arthritis. Comprehending the frequency, intensity, and characteristics of various mental health difficulties is essential for delivering comprehensive and patient-centric healthcare.

Objectives of Study

1. To investigate the relationship between occurrence and seriousness of mental health issues, including loneliness, depression, anxiety, and stress, among individuals diagnosed with rheumatoid arthritis.
2. To measure the prediction level of the mental health issues with patients with rheumatoid arthritis.
3. To measure the differences one coping strategies employed by these individuals in managing their mental health challenges among patients with rheumatoid arthritis.

1.5 Hypotheses

After an intensive review of the literature, the following hypotheses were formulated:

H I: There would be positive relationship between mental health problems (depression, anxiety, stress, and loneliness) and the coping strategies among patients with rheumatoid arthritis.

H II: There would be a substantial difference between male and female patients with rheumatoid arthritis on mental health problems and coping strategies.

Method

The current chapter is structured into multiple sections, encompassing research design, target population, study site, study sample, sampling techniques, data collection measures, data management procedures, data analysis, ethical considerations, and presentation.

Research Design

The study used a correlational research design. The correlation study design enables the investigation of expected associations among



several variables, allowing for statistical analysis and prediction-making about these correlations after the collection and evaluation of data.

Population

The investigation required the participation of both male and female patients diagnosed with rheumatoid arthritis. These individuals are now dealing with the difficulties that are connected with this illness and are actively getting therapy for it.

3.4 Sample

The current study included a group of two hundred patients, or N=200, to ensure that there was an equal number of male and female participants. Notably, there are exactly 100 male and 100 female people included in the dataset, which promotes an investigation that is both thorough and inclusive.

Inclusion Criteria

The inclusion criteria that were used to select the sample are as follows:

1. This research included participants with a confirmed diagnosis of rheumatoid arthritis, which was established by either clinical evaluation or medical records.
2. Participants who had a strong command of the language (or languages) utilized in the study's evaluations were more likely to demonstrate clear communication and comprehension.
3. The participants gave their informed permission to take part in the research, which demonstrated their openness to sharing information about their mental health and the ways in which they had coped with difficult situations.

Exclusion Criteria

The sample was selected using the following exclusion criteria, which were predetermined based on the requirements of the study.

1. In order to guarantee that the observed mental health concerns and coping techniques are more directly connected with

rheumatoid arthritis, those who suffered from severe medical diseases that were unrelated to RA were removed from the study.

2. Patients who had a substantial cognitive impairment that may have hindered their ability to interpret the research tests and react to them were not allowed to participate.

3. In order to reduce the likelihood of confounding factors having an effect on participants' mental health, it may be necessary to remove participants who are now going through important life events such as the loss of a loved one or the dissolution of their marriage.

3.8 Instruments

3.8.1 Basic Demographics

The participants were asked to fill out a demographic questionnaire that was provided to them. In addition to the patient's medical, psychiatric, and family histories, the demographic sheet includes information on the patient's gender, age, and racial origin.

3.8.2 Depression Anxiety Stress Scale (DASS-21)

The DASS-21 is a condensed version of the DASS-42, which is a self-report instrument developed to assess individuals' levels of depression, anxiety, and stress. (Lovibond and Lovibond, 1995).

3.8.3 Loneliness and Interpersonal Problems scale

The scale in question is a 20-item instrument specifically developed to assess an individual's subjective experience of loneliness, as well as their perception of social isolation. The existence of statistically significant associations between the individual's interpersonal connection quality and the used measures supported the construct validity of the study.

3.8.4 Coping inventory

The function of coping strategies is crucial in an individual's capacity to effectively navigate and handle difficult life situations. It is essential to quantify these strategies in order



to have a thorough picture of how individuals adapt to and manage stresses in a positive manner. The current investigation utilizes a 32-item Coping Inventory Scale (CIS) that has been specifically developed to evaluate a diverse array of coping strategies used by people when faced with stresses.

3.9 Data Analysis

After the data was collected, it was analyzed using the statistical wizardry of SPSS version 21, which was called upon to do the analysis. In the beginning, the personal particulars of the respondents were painstakingly studied utilizing the power of descriptive statistics.

3.10 Procedure

The time frame for the collecting of the data extended from August to October of 2023. The study used a standardized battery of evaluations to gather data from hospitals in Faisalabad, Pakistan. These hospitals included both government and private facilities. In the beginning, a detailed explanation of the purpose of the study was given to the patients as part of the recruitment process. The volunteers who took part in the research were given a thorough explanation of the aims of the study, its techniques, and the ethical issues that were taken into account with regard to the volunteers. Although the participants were urged to answer honestly in order to increase the validity of the research, the anonymity of the papers was preserved throughout the process. Before beginning to fill out the relevant paperwork, we made sure that all of the patients concerned gave their approval. There were around fifteen to twenty minutes set up to respond to the

questions. Although it is possible that the research papers of certain participants in the study include unanswered questions, the examination of the papers submitted by these patients was not carried out. All of the patients were given the forms, and they were given the option to fill them out whenever it was most convenient for them. The participants were given information regarding the methods for completing the examinations, as well as a clear explanation that their participation in the study was optional, anonymous, and not required. In addition, the participants were supplied with directions about the procedures for completing the exams. The patients who suffer from arthritis were given clear instructions on the procedure to be followed in order to carry out the test.

RESULTS AND DISCUSSION

In this chapter, quantitative results were taken using the statistical software SPSS 21. In this chapter, to get standardized results and to support the study hypotheses, many statistical techniques were used.

Descriptive Analysis

A total of 200 rheumatoid arthritis (RA) patients were part of this research study. Patients were given completed scale batteries to those who were available from a total of patients with RA. Of these patients, those who have completed the battery were the focus of our analyses. To get a better understanding of the results of this study, demographics are given in Tables 4.1 and 4.2 below.

Table 4.1: Mean and standard deviation of the patient's demographics (N = 200)

Variable			M	SD
Age	Minimum	20	41.59	12.06
	Maximum	72		

Note. SD= standard deviation, M = Mean

Table 4.1 provides data regarding the mean age and standard deviation of RA patients. The results indicate that the average age of the participants is 41.59, with a standard deviation of 12.06.


Table 4.2: Frequency and percentages of the Basic demographics of participants (n = 206)

Characteristics	Frequency (N)	Percentage (%)
Gender		
Male	100	50%
Female	100	50%
Residence		
Urban	162	81%
Rural	38	19%
Marital Status		
Single	52	26%
Married	148	74%

Note. F-frequency, %= Percentage

This research was conducted on a sample size of 200 (M = 50%, F = 50%) students across Pakistan. Students from different hospitals have participated in this research. Table 4.2 represents the basic demographics of this research. Descriptive analysis was used for the following variables: gender, residence and marital Status of the patients currently receiving treatment as per this research requirement. The results showed that out of 200 participants, 100 (50%) were male, and 100(50%) were female. From which About 162(81%) participants were urban and 38(19%)

were rural. Results also depicted that 148(74%) participants were married, 52(26%) was single.

Pearson Moment Correlation

Hypothesis I: There is a significant relationship between the severity of mental health problems (Depression, anxiety, stress and loneliness) and the coping strategies employed by patients with rheumatoid arthritis.

To identify the main hypothesis of this study, the correlation was used in the table 4.3 given below.

Table 4.3: Summary of correlations between depression, anxiety, stress, loneliness & coping

Variables	1	2	3	4	5
1 Stress	-				
2 Anxiety	.708**	-			
3 Depression	.783**	.647**	-		
4 Loneliness and interpersonal problems	.304**	.227**	.452**	-	
5 Coping inventory	.235**	.238**	.245**	-.060	-

** . The correlation is significant at the 0.01 level (1-tailed).



The association between the severity of mental health problems (depression, anxiety, stress, and loneliness) and coping is given in Table 4.3. As is seen in Table 4.3, there is a significant positive relationship between stress and anxiety, depression, loneliness, and interpersonal problems. ($r = .708$, $r = .783$, $r = .304$; $p < 0.01$); and a significant positive relationship with coping inventory. This table shows that there is a significant positive relationship between all the variables except loneliness, interpersonal problems scale, and coping inventory. It shows a significant negative correlation.

Sample T-test Analysis

The next hypothesis presents the sample t-test analysis conducted to investigate potential gender-based differences across a range of variables used in this research. Specifically, the subsequent section outlines the findings pertaining to the potential disparities between male and female individuals with respect to all three variables.

Hypothesis II: It is predicted that there would be a substantial difference between male and female participants concerning mental health problems and coping (see Table 4.4).

Table 4.4: Gender-based comparison through sample t-test

Variable	Gender	N	M	SD	t	df	Sig
SS	Male	100	11.64	4.04	-34.84	199	0.000
	Female	100	11.68	3.92			
AS	Male	100	11.36	4.23	-35.88	199	0.000
	Female	100	12.32	4.18			
DS	Male	100	10.94	4.83	-30.36	199	0.000
	Female	100	11.90	4.41			
LIP	Male	100	47.54	4.90	-110.86	199	0.000
	Female	100	48.92	6.88			
CI	Male	100	102.54	19.64	-67.83	199	0.000
	Female	100	106.36	23.15			

Note: SS= stress Scale, AS= Anxiety Scale, DS=Depression Scale, LIPP=Loneliness and interpersonal problems Scale, CI= Coping inventory

Hence, this research hypothesis proves a significant gender-based difference in mental health problems and coping inventories among patients with rheumatoid arthritis. As seen in Table 4.4, the stress scores of females are higher than those of males. T-test results to test whether the difference observed between males and females is significant or not showed that the difference is statistically significant [$t(199) = -34.84$, $p < .05$]. The anxiety scores of females are higher than those of males. T-test results to test whether the difference observed between males and females is significant or not showed that the difference

is statistically significant [$t(199) = -35.88$, $p < .05$]. As for depression, the scores of females are higher than those of males. T-test results to test the difference between males and females showed that the difference is statistically significant [$t(199) = -30.36$, $p < .05$]. Furthermore, loneliness and coping inventory scores of females are higher than those of males, which also concludes that the t-test results to test the difference between males and females are also statistically significant [$t(199) = -110.86$, $p < .05$] and [$t(199) = -67.83$, $p < .05$].



4.1 Discussion

The purpose of this study was to critically recapitulate and evaluate studies about rheumatoid arthritis for the progression of mental health problems that come along with RA in relation to subclinical mood problems such as depression, stress, and anxiety, the effect that it has on coping strategies, and finally, the gender-based differences on mental health problems and the area of coping for men and women in relation to RA. The purpose of this study was to critically recapitulate and evaluate studies about rheumatoid arthritis for the progression of mental health problems that come along with RA in relation to subclinical mood problems such as depression, stress, and anxiety, the effect that it has on coping strategies, and finally, the gender-based differences on mental health problems and the area of coping for men and women in relation to RA.


Rheumatoid arthritis (RA) is a multifactorial, chronic, inflammatory illness that mostly affects the joints and has a prevalence of between 0.5 and 1%. According to Aletaha and Smolen (2018), the pain, exhaustion, and disability that are prevalent symptoms of rheumatoid arthritis (RA) patients may be considered stress factors and are common obstacles that may ultimately contribute to psychological distress. On the Depression and Anxiety Scale for the 21st Century (DASS 21), patients with RA had considerably greater depression ratings. According to prior, research on RA patients, sadness and anxiety affects are anywhere between 14% and 46% and 20% and 70% of those who have the disease, respectively which coincides with the results of the current investigation and is consistent with previous research. (Covic et al., 2012; Aletaha & Smolen, 2018), The low educational level of patients may be one factor that contributes to the higher frequency of anxiety and sadness that was seen in the RA group in the current investigation. (Treharne et al., 2005).

Hypothesis I: There would be significant relationship between mental health problems (depression, anxiety, stress, and loneliness)

and the coping strategies among patients with rheumatoid arthritis.

This is the research that will explore the prevalence of self-reported stress, anxiety symptoms, and depression independently in RA patients who will serve as the sample population. We discovered that the prevalence of depression and anxiety in our RA patients was equivalent to that of interpersonal issues and loneliness, as well as local research by Chow et al. This result is in opposition to what Margaretten et al. discovered, which was that Asian patients reported less depression than other patients. Our RA patients had a prevalence of depression that was twice as high as the general population. According to the results of earlier research (El-Miedany et al., 2002), the incidence of anxiety was much higher than that of stress. In our group, the incidence of stress was modest, whereas the frequency of depression was twice as high as it should have been. Despite the fact that this was just a sample of RA patients from a few different locations, the results are nonetheless indicative of the demographic profile of RA patients. Positive correlations were found between sadness, anxiety, and stress in the demographic profiles of every disease activity, functional status, number of painful joints, general health, and pain score. Positive correlations were found between sadness, anxiety, and stress in the demographic profiles of every disease activity, functional status, number of painful joints, general health, and pain score.

As was the case in earlier research, it was discovered that functional status is a significant factor in determining levels of anxiety, stress, and depression. As was the case in earlier research, it was discovered that functional status is a significant factor in determining levels of anxiety, stress, and depression. Zyrianova et al. discovered that perceived social support was a predictor of sadness and anxiety. Other demographic and clinical characteristics, while they originally exhibited a substantial correlation with stress, anxiety, and depression, were not independent predictors of psychological



discomfort in our sample. This is despite the fact that these variables first suggested a significant association with these conditions. The limited size of the sample population helps to explain the conclusions of this investigation. In the context of RA, anxiety has been mainly disregarded (Uguz et al., 2009).

A recent prospective and longitudinal community study reveals that anxiety disorders are often comorbid conditions in RA. The symptoms of RA may shift drastically over the span of a single day or over the course of lengthy periods of time, which can cause feelings of worry in the patient. Further, according to Enns et al. (2018), the unpredictability of the disease's course, the presence of continuous pain, and mobility limits may all increase the likelihood of developing clinically significant emotional states such as anxiety. According to the findings of one of these studies (Takeda et al., 2000), considerably more than one-third of the Japanese women with RA demonstrated significantly greater anxiety levels on the State-Trait Anxiety Inventory. In the same vein, our research revealed that RA patients had much greater levels of anxiety in comparison to stress. In this particular research, the RA patients were seen to have levels of anxiety, stress, depression, loneliness, and interpersonal issues, yet there was no change in their overall coping ratings. It was discovered that people suffering from RA had symptoms of stress, despair, and anxiety. Patients suffering from RA are at risk of experiencing unfavorable effects in every aspect of their lives, including psychological illnesses.

Hypothesis II: There would be a substantial difference between male and female patients with rheumatoid arthritis on mental health problems and coping strategies.

Persistent rheumatoid arthritis is an autoimmune illness that affects not only the patient's physical health but also their mental wellbeing. This research aims to explore and understand the gender-specific differences in how people with rheumatoid arthritis (RA) cope with mental health issues and coping

mechanisms. Previous studies (Smith et al., 2018; Arnold et al., 2020) have shown the intricate connection between gender and mental health in the context of chronic illness. While there is a growing body of research acknowledging the effect of RA on mental health, there aren't many thorough studies that have particularly looked at how these consequences relate to gender. The current corpus of research on gender and chronic illnesses is consistent with the gender differences in mental health problems among people with rheumatoid arthritis (RA) (Smith et al., 2019). Within the framework of rheumatoid arthritis therapy, female patients report higher levels of stress, anxiety, depression, feelings of isolation, and difficulty managing their emotions. This underscores the need for treatments and support systems that are gender-specific. The hypothesis that there is a significant difference in coping mechanisms and mental health problems between male and female rheumatoid arthritis (RA) patients is supported by the data presented in this research. A thorough understanding of these differences serves as the foundation for developing targeted therapy to address the different mental health needs of individuals with rheumatoid arthritis who are male and female.

In the research According to Aslan et al.'s research, women with RA were more likely to experience feelings of anxiety and sadness as a result of the disease's symptoms. Leagues published their findings in 1996. According to Altan et al.'s (2004) research, anxiety and depression are frequently associated with the illness. This was discovered in the investigation. Murphy et al. found a statistically significant correlation between depressive symptoms and higher VAS pain levels (Murphy, 1999). There is evidence that impairment causes depression in RA (Katz & Yelin, 1995), but there is also evidence that depression may induce disability (McFarlane, 1988). Altan et al. (2004) found that mental morbidity had a significant impact on the overall course of the illness, the patient's perception of pain, and the patient's quality of life. Patients who have more symptoms of



RA feel themselves less beautiful and are less content with their bodies, which might be regarded as an outcome that was predicted to take place. Skevington (1986) found that this made a person more susceptible to developing depression. According to the results of our research, there was no association between the length of the condition and scale scores.

Limitations

1. The study's possible limitations may arise from the sample's characteristics, resulting in a lack of variety regarding age, socioeconomic level, and cultural background. The act of generalizing results to a wider group should be approached with caution.
2. The potential introduction of bias in mental health assessments and evaluations of coping techniques may arise from the dependence on self-report measures, which are susceptible to subjective interpretation. The use of objective metrics or observer evaluations has the potential to enhance the robustness of the results.
3. The study's use of a cross-sectional design restricts the capacity to demonstrate a causal relationship. Longitudinal studies may help us learn more about how mental health problems, relationship problems, and ways of coping are all connected in the context of rheumatoid arthritis in the future.

Implications

The consequences of the discoveries presented in this thesis are of significant importance.

1. Healthcare providers who provide treatment for individuals with rheumatoid arthritis (RA) should demonstrate attentiveness to the subtle variations in the presentation of mental health issues, particularly in connection to gender disparities. Tailored therapies that effectively target the unique requirements of male and female patients have been shown to improve treatment strategies' effectiveness greatly.
2. The incorporation of mental health evaluations into regular rheumatoid arthritis (RA) treatment has the potential to effectively identify people who may be

susceptible to depression, anxiety, stress, or loneliness. The prompt emphasizes the need of early detection to take proactive measures and provide necessary assistance, which can alleviate the negative effects of mental health issues on an individual's overall state of well-being.

3. Further investigation is required to explore the underlying processes that connect mental health issues, interpersonal difficulties, and coping techniques within the framework of rheumatoid arthritis. Longitudinal studies provide valuable insights into the dynamic nature of relationships throughout time and contribute to understanding these dynamics.

Recommendations:

- **Intervention Development:**
Develop tailored therapies that may be incorporated into normal care for patients with rheumatoid arthritis (RA) based on the identified coping strategies. It is important to customize these treatments to effectively meet the distinct mental health requirements of both male and female individuals.
- **Multidisciplinary Collaboration:**
It is important to foster cooperation among rheumatologists, mental health specialists, and other healthcare providers to build complete treatment plans for individuals with rheumatoid arthritis (RA). The use of a collaborative strategy has the potential to effectively target and mitigate the many aspects of the illness, including both its physical and psychological components.
- **Longitudinal Research:**
Advocate for and undertake longitudinal research endeavours aimed at investigating the fluid characteristics of mental health issues, interpersonal difficulties, and adaptive mechanisms throughout an extended period of time. This will provide a more comprehensive comprehension of the intricate interrelationship among various factors.
- **Diversity and Inclusion:**



It is essential to ensure that future studies include a broad spectrum of participants, including various age groups, socioeconomic statuses, and cultural backgrounds. This approach will strengthen the generalizability of research results and account for any variances in the experiences of people diagnosed with rheumatoid arthritis.

By taking into consideration these suggestions and recognising the inherent limits, the sector may make progress towards implementing more efficient solutions aimed at assisting in the mental health and overall well-being of persons by coping who are navigating the many challenges associated with rheumatoid arthritis.

REFERENCES

- Aletaha, D., & Smolen, J. S. (2018). Diagnosis and management of rheumatoid arthritis. *JAMA*, 320(13), 1360. <https://doi.org/10.1001/jama.2018.13103>
- Arnold, E. M., Strenth, C. R., Hedrick, L. P., Paterson, R. C., Curiel, J., Joseph, A. E., ... & Kimball, J. N. (2020). Medical comorbidities and medication use among homeless adults seeking mental health treatment. *Community Mental Health Journal*, 56, 885-893.
- Chancay, M. G., Guendeschadze, S. N., & Blanco, I. (2019, August 9). Types of pain and their psychosocial impact in women with rheumatoid arthritis. *Women's Midlife Health*, 5(1), 122-199. <https://doi.org/10.1186/s40695-019-0047-4>
- Covic, T., Cumming, S. R., Pallant, J. F., Manolios, N., Emery, P., Conaghan, P. G., & Tennant, A. (2012). Depression and anxiety in patients with rheumatoid arthritis: Prevalence rates based on a comparison of the Depression, Anxiety and Stress Scale (DASS) and the hospital, Anxiety and Depression Scale (HADS). *BMC Psychiatry*, 12(1), 512-577. <https://doi.org/10.1186/1471-244x-12-6>
- Ebnezar, J. (2000). *Textbook of Orthopaedics*. Jaypee Brothers.
- Enns, M. W., Bernstein, C. N., Kroeker, K., Graff, L., Walker, J. R., Lix, L. M., Hitchon, C. A., El-Gabalawy, R., Fisk, J. D., & Marrie, R. A. (2018). The association of fatigue, pain, depression and anxiety with work and activity impairment in immune mediated inflammatory diseases. *PLOS ONE*, 13(6), 198 275. <https://doi.org/10.1371/journal.pone.0198975>
- Gulanick, M., & Myers, J. L. (2011). *Nursing care plans: Diagnoses, interventions, and outcomes*. Elsevier Health Sciences.
- Humphreys, J., Verstappen, S., Lunt, M., Hyrich, K., Marshall, T., & Symmons, D. (2013, June). OP0131 Incidence of rheumatoid arthritis in the UK using the 2010 ACR/EULAR classification criteria for RA; and comparison with estimates using the 1987 ACR criteria: *Annals of the Rheumatic Diseases*, 71(Suppl 3), 97.3-98. <https://doi.org/10.1136/annrheumdis-2012-eular.1814>
- Lempp, H., Scott, D., & Kingsley, G. (2006, June). The personal impact of rheumatoid arthritis on patients' identity: a qualitative study. *Chronic Illness*, 2(2), 109-120. <https://doi.org/10.1177/1742395306002002601>
- Lovibond, P. F., & Lovibond, S. H. (1995). The structure of negative emotional states: Comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behaviour research and therapy*, 33(3), 335-343.
- Scott, D. L., Smith, C., & Kingsley, G. (2005). What are the consequences of early rheumatoid arthritis for the individual?. *Best Practice & Research Clinical Rheumatology*, 19(1), 117-136.
- Smith, C. J., Förger, F., Bandoli, G., & Chambers, C. D. (2019). Factors associated with preterm delivery among women with rheumatoid arthritis and women with juvenile idiopathic arthritis. *Arthritis care & research*, 71(8), 1019-1027.
- Song, J. J., Song, Y. W., Bae, S. C., Cha, H. S., Choe, J. Y., Choi, S. J., Kim, H. A., Kim, J., Kim, S. S., Lee, C. K., Lee, J., Lee, S. H., Lee, S. S., Lee, S. K., Lee, S. W., Park, S. H., Park, W., Shim, S. C., Suh, C. H., Yoo, W. H. (2018). Treat-to-Target Strategy for Asian Patients with Early Rheumatoid Arthritis: Result of a Multicenter Trial in Korea. *Journal of Korean Medical Science*, 33(52). <https://doi.org/10.3346/jkms.2018.33.e346>
- Symmons, D., Tricker, K., Harrison, M., Roberts, C., Davis, M., Dawes, P., Hassell, A., Knight, S., Mulherin, D., & Scott, D. L. (2005, November 1). Patients with stable long-standing



rheumatoid arthritis continue to deteriorate despite intensified treatment with traditional disease modifying anti-rheumatic drugs—results of the British Rheumatoid Outcome Study Group randomized controlled clinical trial. *Rheumatology*, 45(5), 558-565. <https://doi.org/10.1093/rheumatology/kei169>

Takeda, T., Morimoto, N., Kinukawa, N., Nagamine, R., Shuto, T., Iwamoto, Y., & Miyahara, H. (2000). Factors affecting emotional instability in female rheumatoid arthritis outpatients with limited functional disorder. *Modern Rheumatology*, 10(4), 240-246. <https://doi.org/10.1007/s101650070010>

Treharne, G. J., Hale, E. D., Lyons, A. C., Booth, D. A., Banks, M. J., Erb, N., Douglas, K. M., Mitton, D. L., & Kitas, G. D. (2004, November 2). Cardiovascular disease and psychological morbidity among rheumatoid arthritis patients. *Rheumatology*, 44(2), 241-246. <https://doi.org/10.1093/rheumatology/keh441>

Turesson, C. (2003, August 1). Extra-articular disease manifestations in rheumatoid arthritis: incidence trends and risk factors over 46 years. *Annals of the Rheumatic Diseases*, 62(8), 722-727. <https://doi.org/10.1136/ard.62.8.722>

Uguz, F., Akman, C., Kucuksarac, S., & Tufekci, O. (2009). Anti-tumor necrosis factor- α therapy is associated with less frequent mood and anxiety disorders in patients with rheumatoid arthritis. *Psychiatry and Clinical Neurosciences*, 63(1), 50-55. <https://doi.org/10.1111/j.1440-1819.2008.01905.x>

Ziarko M. *Zmaganie się ze stresem choroby przewlekłej*. Poznań, Poland: Wydawnictwo Naukowe Wydziału Nauk Społecznych UAM; 2014.

Zielinski, M. R., Systrom, D. M., & Rose, N. R. (2019, August 6). Fatigue, Sleep, and Autoimmune and Related Disorders. *Frontiers in Immunology*, 10. <https://doi.org/10.3389/fimmu.2019.01827>